(19) World Intellectual Property Organization International Bureau



(43) International Publication Date 3 January 2002 (03.01.2002)

PCT

(10) International Publication Number WO 02/00198 A1

(51) International Patent Classification7:

A61K 9/16

- (21) International Application Number: PCT/GB01/02922
- (22) International Filing Date: 29 June 2001 (29.06.2001)
- (25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

0015981.4

29 June 2000 (29.06.2000) (

- (71) Applicant (for all designated States except US): GLAXO GROUP LIMITED [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): LANCASTER, Robert, William [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). SINGH, Hardev [IN/GB]; GlaxoSmithKline, Temple Hill, Dartford, Kent DA1 5AH (GB). THEOPHILUS, Andrew, Lewis [GB/GB]; GlaxoSmithKline, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB).
- (74) Agent: GIDDINGS, Peter, John: GlaxoSmithKline, Corporate Intellectual Property, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).

- (81) Designated States (national): AE. AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- -- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

02/00198 AJ

(54) Title: PROCESS FOR PREPARING AND HARVESTING CRYSTALLINE PARTICLES

(57) Abstract: The present invention relates to a novel process for preparing and harvesting crystalline particles, particularly particles of therapeutically useful or carrier substances of a size suitable for inhalation therapy.

C 47 C DO 1702722

process for preparing and harvesting crystalline particles

This invention relates to a novel process for preparing crystalline particles, particularly particles of defined particle size distribution, especially particles of therapeutically useful or carrier substances of a size suitable for inhalation therapy.

5

10

15

20

25

Industrial processes for production of many products, particularly pharmaceutical products, require the preparation of pure substances of a defined particle size distribution. Pure substances are frequently prepared by precipitation from solutions of lesser purity. When precipitation takes place relatively slowly (e.g. over a matter of hours), crystals are grown which are frequently of a non-uniform shape and relatively large size.

In the field of inhalation therapy, therapeutic molecules are generally desired of a particle size "suitable for inhalation", which is a term generally taken to indicate an aerodynamic diameter between 1 and 10 μm , especially 1 and 5 μm , particularly 1 and 3 μm . Carrier molecules (such as lactose) for inhaled therapeutic preparations are typically desired of a significantly larger aerodynamic diameter so that they do not penetrate into the upper respiratory tract to the same degree as the active ingredient and an aerodynamic diameter of 100 to 150 μm is generally considered suitable. However this is a generalisation and for some purposes it may well be preferred to use a lower particle size for the carrier, even one comparable to that of the therapeutic substance.

Outside of the inhaled area, modification of the habit and size of crystals is a valuable tool in adjusting and optimising pharmaceutical and biological properties such as flow characteristics, dissolution rate and bioavailability.

10

15

20

25

2

Particles of the desired particle size for inhalation therapy are conventionally prepared by milling or micronisation. These processes, depending on the precise conditions adopted, are capable of generating particle distributions which include fractions having particles with the appropriate size. suitable for preparing particles of the larger size indicated above and micronisation of the smaller size indicated above. However, there are a number of disadvantages associated with milling and micronisation processes including that the fraction having the desired particle size may be relatively small, that there may be generated a significant fraction of particles that are finer than is desired (which may be deleterious e.g. if it affects bioavailability) and that product losses generally may be considerable (e.g. through coating of the machinery). A further property of micronised products is that the surfaces of the particles generated are generally substantially amorphous (i.e. have minimal crystallinity). This may be undesirable when there exists a tendency for the amorphous regions to convert to a more stable crystalline state. Furthermore micronised or milled products may be more susceptible to moisture uptake than crystalline products. Micronisation and milling processes also suffer from the disadvantages that they are relatively energy intensive and require containment and other measures to avoid the risk of dust explosion.

International patent application PCT/GB99/04368 (filed but not published before the priority date of this application) describes a process and apparatus for preparing particles which comprises mixing in the presence of ultrasonic radiation a flowing solution of a substance in a liquid solvent with a flowing liquid antisolvent for said substance. International patent application PCT/GB00/04237 describes a process which comprises admitting a stream of a solution of a substance in a liquid solvent and a stream of liquid antisolvent for said substance tangentially into a cylindrical mixing chamber, consequently causing a

vortex which results in precipitation of crystalline particles. However, the disadvantage with these 2 processes is that particle growth or agglomeration may occur in the course of isolating the particles from the solvent/anti-solvent mixture. We have now invented an improvement to these processes which is less susceptible to the above mentioned disadvantage.

Thus, according to a first aspect of the invention there is provided a process for preparing crystalline particles of a substance which comprises mixing a flowing solution of the substance in a liquid solvent with a flowing liquid antisolvent for said substance in order to generate a suspension of crystalline particles in the solvent/anti-solvent the process further comprises the steps of

- (a) filtering the suspension of crystalline particles in the solvent/anti-solvent mixture in order to remove the solvent/antisolvent mixture;
- (b) washing the filtered particles with anti-solvent;
- (c) resuspending the filtered and washed particles in anti-solvent;
 - (d) cooling the resultant suspension of filtered, washed and resuspended particles in the anti-solvent; and
 - (e) collecting crystalline particles by removal of the antisolvent from the cooled suspension.

20

25

15

5

10

In a first preferred embodiment of the present invention said mixing comprises mixing in a continuous flow cell in the presence of ultrasonic radiation.

In a second preferred embodiment of the present invention said mixing comprises admitting a stream of solution of the substance in a liquid solvent and a stream of liquid antisolvent for said substance tangentially into a cylindrical mixing chamber having an axial outlet port such that said streams are thereby intimately mixed through formation of a vortex and precipitation of crystalline particles of the substance is thereby caused.

Preferably, the solvent will be miscible with the anti-solvent.

Preferably, the suspension of crystalline particles in the solvent/anti-solvent mixture will be filtered using a wide range of suitable filters known to persons skilled in the art. Examples of filters include sinters (e.g. glass sinters), fibre filters (e.g. paper and nitrocellulose filters) and membrane filters. We have found that a particularly advantageous filtration arrangement involves use of a glass fibre microfilter sandwiched between two Whatman paper filters (e.g. Whatman 54 filters). The particle size of the filter will be appropriate for the product collected. It is possible to modify the distribution of particles at the fine end by selecting a filter size which allows fines to pass through the filter. Preferably, the filter will be a filter suitable to retain crystalline particles of between 1 and 10μm, most preferably less than 5μm, especially less than 3μm.

15

10

5

It will be appreciated that the anti-solvent used in washing step (b) and resuspension step (c) does not need to be the same anti-solvent that is used in the original process which generates the crystalline particles. Preferably, however, the anti-solvent used in washing step (b) and resuspension step (c) will be the same anti-solvent as is used in the original process.

20

Preferably, the suspension of crystalline particles obtained in step (d) will be cooled to freezing point. Also preferably, the suspension of crystalline particles obtained in step (a) will be cooled to freezing point using a solid carbon dioxide cooling bath containing a suitable solvent eg. acetone, IMS or methanol.

25

Where possible, the antisolvent will preferably be water. Preferably, in step (d) the removal of the antisolvent from the cooled suspension is achieved by freeze drying.

The process of the present invention has the advantage of maintaining the original particle diameter of the particles of substance achieved by crystallisation. Conventional collection techniques involve further incubation of the particles in the solvent/antisolvent mixture which may result in undesirable effects such as crystal growth. Wherein the particles are prepared for inhalation therapy, crystal growth is disadvantageous because the particles may grow to a diameter such that they may not be effectively delivered to the lower respiratory airways.

10

15

20

5

The advantages that the invention may possess include the fact that the process may be performed in a continuous manner without requirements for batch processing, that the process may be scaled up with relative ease and that the process is capable of producing particle size distributions of very high uniformity index.

Surprisingly, the present invention provides processes for removing the solvent from the solvent/antisolvent mixture in order to prevent crystal growth, and as demonstrated in the Examples, also results in particles with more refined particle sizes than achieved with conventional harvesting techniques. Furthermore, when the antisolvent is water, once the solvent has been removed from the solvent/antisolvent mixture (by either procedure) and the mixture is cooled to freezing point, the freeze drying step ensures that the water molecules sublime from the mixture leaving only particles containing the desired substance(s).

25 substance(s)

The process of the present invention is particularly suitable for preparing particles of substances which are pharmaceutical or carrier substances suitable for inhalation therapy.

Substances suitable for inhalation therapy include substances applied topically to the lung and nose.

Examples of pharmaceutical substances suitable for inhalation therapy include 5 analgesics, e.g., codeine, dihydromorphine, ergotamine, fentanyl or morphine; anginal preparations, e.g., diltiazem; antiallergics, e.g., cromoglycate, ketotifen or nedocromil; antiinfectives e.g., cephalosporins, penicillins, streptomycin, sulphonamides, tetracyclines and pentamidine; antihistamines, e.g., methapyrilene; anti-inflammatories, e.g., beclomethasone (eg. as the 10 dipropionate), fluticasone (eg. as the propionate), flunisolide, budesonide, rofleponide, mometasone (e.g. as the furoate) or triamcinolone (e.g. as the acetonide); antitussives, e.g., noscapine; bronchodilators, e.g., albuterol (eg. as the sulphate), salmeterol (eg. as the xinafoate), ephedrine, adrenaline, fenoterol (eq. as the hydrobromide), formoterol (e.g. as the fumarate), isoprenaline, 15 metaproterenol, phenylephrine, phenylpropanolamine, pirbuterol (eg. as the acetate), reproterol (eg. as the hydrochloride), rimiterol, terbutaline (eg. as the sulphate), isoetharine, tulobuterol or (-)-4-amino-3,5-dichloro- α -[[[6-[2-(2pyridinyl)ethoxyl hexyllmethyll benzenemethanol; diuretics, e.g., amiloride; anticholinergics, e.g., ipratropium (e.g. as the bromide), tiotropium, atropine or 20 oxitropium; hormones, e.g., cortisone, hydrocortisone or prednisolone; xanthines, e.g., aminophylline, choline theophyllinate, lysine theophyllinate or theophylline; therapeutic proteins and peptides, e.g., insulin or glucagon; and salts, esters and solvates of any of the above. Other examples include 4hydroxy-7-[2-[[2-[[3-(2-phenylethoxy)propyl]sulfonyl]ethyl] amino]ethyl-2(3H)-25 benzothiazolone or butixicort and salts and solvates thereof. Another example of a pharmaceutical substance suitable for inhalation therapy is 6α , 9α -difluoro-11 β -hydroxy-16 α -methyl-3-oxo-17 α -propionyloxy-androsta-1.4-diene-17β-carbothioic acid S-(2-oxo-tetrahydro-furan-3-yl) ester or a solvate thereof (which compound is especially suitable for administration by the nasal route).

Other examples of pharmaceutical substances suitable for inhalation therapy which are of particular interest are:

5 (2R,3R,4S,5R)-2-[6-Amino-2-(1S-hydroxymethyl-2-phenyl-ethylamino)-purin-9-yl]-5-(2-ethyl-2H-tetrazol-5-yl)-tetrahydro-furan-3,4-diol or a salt thereof (eg. the maleate salt); and

(2S)-3-[4-({[4-(Aminocarbonyl)-1-piperidinyl]carbonyl}oxy)phenyl]-2-[((2S)-4-methyl-2-{[2-(2-methylphenoxy)acetyl]amino}pentanoyl)amino] propanoic acid or a salt thereof (eg. as free acid or potassium salt).

Examples of other pharmaceutical substances for which the process according to the invention is useful include compounds to be administered orally such as 2(S)-(2-benzoyl-phenylamino)-3-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid, 2,6-diamino-3-(2,3,5-trichlorophenyl)pyrazine and naratriptan (eg. as hydrochloride) and other 5HT-1 agonists such as sumatriptan (eg. as succinate). Another compound of interest is (S)-[2-(1-iminoethylamino)ethyl]-L-homocysteine or a salt or racemate thereof (eg. preferably the 2- isomer).

20

BNEDOCID: -INO

10

Pharmaceutical substances as described above include asymmetric molecules which may exist as mixtures of optical isomers (e.g. as racemates) or as purified single enantiomers.

Pharmaceutical substances of particular interest include fluticasone, beclomethasone, salmeterol, salbutamol or an ester, salt or solvate thereof. The substance of most interest is salmeterol xinafoate (including the racemate or the purified r- or s- enantiomers). Fluticasone propionate is also of particular interest.

10

15

20

25

Examples of carrier substances include lactose.

The solvent and antisolvent liquids will be selected so as to be appropriate for the substance. Preferably, they are readily miscible in the proportions employed. Suitable combinations of solvent/antisolvent include acetone/water, ethanol/IPA, methanol/IPA, methanol/water and reciprocal pairs. Methanol/IPE is also a suitable pairing.

For generation of small particles by the process according to the invention, it is preferred that the difference between the dissolution properties of the solvent and anti-solvent be as great as possible. For reasons of industrial efficiency (particularly in order to reduce the throughput volumes of liquid) it is preferred to use concentrations of substance in solvent which are as high as possible. Nevertheless the solutions must be stable and not prone to crystallisation before discharge into the continuous flow cell. With this end in mind, it may be preferred to use the solution of the substance in the solvent at elevated temperature. It may also be preferable to cool the anti-solvent.

In order to prevent premature precipitation of the dissolved substance in the lines it will generally be desired to prime the apparatus by first pumping it with solvent. It may be preferred to prime the apparatus by pumping it with heated solvent, particularly when the dissolved substance is close to its solubility limit.

When the substance is fluticasone propionate we prefer the solvent to be acetone and the anti-solvent to be water.

When the substance is salmeterol xinafoate we prefer the solvent to be methanol or acetone (more preferably methanol) and the anti-solvent to be water.

When the substance is salbutamol sulphate we prefer the solvent to be water and the anti-solvent to be IMS.

When the substance is beclomethasone dipropionate we prefer the solvent to be IMS and the anti-solvent to be water.

When the substance is lactose we prefer the solvent to be water and the antisolvent to be ethanol.

When the substance is budesonide, we prefer the solvent to be methanol and the anti-solvent to be water.

When the substance is formoterol fumarate or terbutaline sulphate we prefer the solvent to be methanol or acetone and the anti-solvent to be water.

When the substance is 2,6-diamino-3-(2,3,5-trichlorophenyl)pyrazine we prefer the solvent to be methanol and the anti-solvent to be water.

15

20

25

When the substance is 2(S)-(2-benzoyl-phenylamino)-3-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid we prefer the solvent to be acetone and the anti-solvent to be water.

When the substance is naratriptan hydrochloride we prefer the solvent to be methanol and the antisolvent to be IPE.

When the substance is 6α , 9α -difluoro- 11β -hydroxy- 16α -methyl-3-oxo- 17α -propionyloxy-androsta-1,4-diene- 17β -carbothioic acid S-(2-oxo-tetrahydro-furan-3-yl) ester we prefer the solvent to be acetone and the anti-solvent to be water.

We have found that the method according to the invention is suitable for producing populations of mixtures when the substance is a mixture of substances. When the substance is a mixture the method has particular advantages since it is capable of producing mixtures of crystalline particles of very high homogeneity without the need for any blending step. When the substance is a mixture the solvent and anti-solvent will have to be appropriate for all components of the mixture. Differential solubilities in the recrystalline mixture tend to result in the output proportions of the mixture differing from the

10

15

20

25

initial proportions in solution in the solvent and so appropriate adjustment of the input proportions to achieve the desired output proportions may be necessary.

The method according to the invention is particularly suitable for producing mixtures of crystalline particles of salmeterol and fluticasone or salts and esters thereof e.g. salmeterol xinafoate and fluticasone propionate. The preferred solvent is acetone. The preferred anti-solvent is water. Recrystallisation from acetone using water as anti-solvent tends to cause an increase in the ratio of salmeterol xinafoate to fluticasone propionate relative to their proportion in solution in acetone. The method is also expected to be suitable for producing mixtures of crystalline particles of formoterol and budesonide or salts and esters thereof e.g. formoterol fumarate and budesonide.

As a further aspect of the invention we provide a population of particles obtainable by a process according to the invention.

Particles of pharmaceutical or carrier substances may be obtained which are suitable for use in a pharmaceutical composition for inhalation therapy, such as dry powder composition (whether containing pure drug, or drug mixed with a carrier such as lactose) or a pressurised liquid formulation (e.g. a formulation comprising a hydrofluoroalkane (HFA) propellant such as HFA134a or HFA227 or a mixture thereof).

Pressurised liquid formulations suitable for metered-dose inhalers will be retained in canisters, typically aluminium canisters (which may be plastics lined) which are provided with a metering valve of appropriate metering volume.

It will be appreciated that references to inhalation therapy also extend to administration of pharmaceutical compositions via the nasal route. Formulations

suitable for nasal delivery include pressurised (e.g. HFA containing) formulations and non pressurised (e.g. aqueous) formulations which may be metered by the delivery device adapted for administration to the nose.

We also provide a pharmaceutical composition comprising a population of particles prepared according to the invention.

Apparatus suitable for use in the present invention is illustrated by reference to Figure 1 in which mixing chamber 1 is provided with first inlet port 2 connected to first reservoir 3 containing substance dissolved in solvent and second inlet port 4 connected to second reservoir 5 containing anti-solvent. Pumps 6 and 7 deliver liquid from reservoirs 3 and 5 to mixing chamber 1 at a controlled rate. An ultrasound probe 8 is located in the vicinity of, and just above, inlet port 2. When pumps 6 and 7 are in operation, liquids from reservoirs 3 and 5 are delivered to mixing chamber 1 and are mixed with the aid of magnetic stirrer 9. Liquid containing the particles of substance thus generated flows out of the mixing chamber via exit port 10. The solvent within this flowing suspension is then removed using a filter 11 according to the present invention.

20 Brief description of the drawings.

Figure 1: Example apparatus according to the invention

The present invention may be illustrated by the following non-limiting Example:

25 Examples

10

15

Example 1: Distributions of particles of crystalline fluticasone propionate

The drug substance (fluticasone propionate) (1wt) was dissolved in hot acetone
(15vol) and then allowed to cool to ambient temperature (20°C). A flow cell was

then charged with a 4:1 mixture of water and acetone respectively. Pump 1 (containing the fluticasone propionate in acetone) was set at a flow rate of 20ml/min. Pump 2 (containing water chilled to 3-5°C) was set at a flow rate of 80ml/min. A magnetic stirrer bar was placed in the flow cell. The tip of a sono-probe was positioned above the inlet of Pump 1 and set to deliver 70-75 watts of power. When the ultrasound probe, both pumps and the magnetic stirrer were turned on, rapid onset of crystallisation occurred.

5

10

15

20

The resultant crystalline suspension was then collected and simultaneously filtered on a filter funnel fitted with GF/C glass microfibre filter sandwiched between 2 Whatman No. 54 filter papers. The damp filter cake was then washed with water and then resuspended in further demineralised water to prepare a 10% w/w slurry. The slurry was then rapidly frozen by immersing the flask containing the slurry in a solid carbon dioxide cooling bath containing acetone, to give an even coating of ice containing particles of fluticasone propionate. The mixture was then freeze dried *in vacuo* for 14-18 hours to give a fine white powder containing particles of inhalable quality.

The contents of the above mentioned patent application is herein incorporated by reference.

CLAIMS

5

- 1. A process for preparing crystalline particles of a substance which comprises mixing a flowing solution of the substance in a liquid solvent with a flowing liquid antisolvent for said substance in order to generate a suspension of crystalline particles in the solvent/anti-solvent the process further comprises the steps of
- (a) filtering the suspension of crystalline particles in the solvent/anti-solvent mixture in order to remove the solvent/antisolvent mixture;
- 10 (b) washing the filtered particles with anti-solvent;
 - (c) resuspending the filtered and washed particles in anti-solvent;
 - (d) cooling the resultant suspension of filtered, washed and resuspended particles in the anti-solvent; and
- (e) collecting crystalline particles by removal of the antisolvent from the cooled suspension.
 - 2. A process according to claim 1 wherein said mixing comprises mixing in a continuous flow cell in the presence of ultrasonic radiation.
- 3. A process according to claim 1 wherein said mixing comprises admitting a stream of solution of the substance in a liquid solvent and a stream of liquid antisolvent for said substance tangentially into a cylindrical mixing chamber having an axial outlet port such that said streams are thereby intimately mixed through formation of a vortex and precipitation of crystalline particles of the substance is thereby caused.
 - 4. A process according to any one of claims 1 to 3 wherein the solvent is miscible with the anti-solvent.

PCT/GB01/02922

5. A process according to any one of claims 1 to 4 wherein the suspension of crystalline particles in the solvent/anti-solvent mixture will be filtered using a filter which is suitable to retain crystalline particles of between 1 and $10\mu m$.

5

- 6. A process according to claim 5 wherein the filter is suitable to retain crystalline particles of less than $5\mu m$.
- 7. A process according to claim 6 wherein the filter is suitable to retain
 10 crystalline particles of less than 3μm.
 - 8. A process according to any one of claims 1 to 7 wherein the antisolvent used in washing step (b) and resuspension step (c) is the same antisolvent as is used in the original process which generates the crystalline particles.
 - 9. A process according to any one of claims 1 to 8 wherein the suspension of crystalline particles obtained in step (d) are cooled to freezing point.

20

15

10. A process according to any one of claims 1 to 9 wherein the suspension of crystalline particles obtained in step (a) are cooled to freezing point using a solid carbon dioxide cooling bath containing a suitable solvent eg. acetone, IMS or methanol.


25

11. A process according to any one of claims 1 to 10 wherein the antisolvent is water.

- 12. A process according to any one of claims 1 to 11 wherein in step (d) the removal of the antisolvent from the cooled suspension is achieved by freeze drying.
- 5 13. A process according to any one of claims 1 to 12 wherein the process prepares particles of substances which are pharmaceutical or carrier substances suitable for inhalation therapy.
- 14. A process according to claim 13 wherein the substance is fluticasone,
 10 beclomethasone, salmeterol, salbutamol or an ester, salt or solvate thereof.
 - 15. A process according to claim 13 wherein the substance is lactose.
- 16. A process according to claim 13 wherein the substance is 6α, 9α 15 difluoro-11β-hydroxy-16α-methyl-3-oxo-17α-propionyloxy-androsta-1,4-diene 17β-carbothioic acid S-(2-oxo-tetrahydro-furan-3-yl) ester.
 - 17. A process according to claim 14 wherein the substance is fluticasone propionate.
 - 18. A process according to claim 14 wherein the substance is salmeterol xinafoate.
- 19. A process according to any one of claims 1, 2, 3 or 11 wherein the substance is a mixture.

20. A process according to claim 19 wherein the substance is a mixture of fluticasone propionate and salmeterol xinafoate.

- 21. A process according to any one of claims 1 to 12 wherein the process prepares particles of substances which may be administered orally.
- 22. A process according to claim 21 wherein the substance is 2(S)-(2-benzoyl-phenylamino)-3-{4-[2-(5-methyl-2-phenyl-oxazol-4-yl)-ethoxy]-phenyl}-propionic acid or 2,6-diamino-3-(2,3,5-trichlorophenyl)pyrazine.
 - 23. A process according to claim 21 wherein the substance is naratriptan hydrochloride.
 - A population of particles obtainable by a process according to any one of claims 1 to 23.
- 25. A pharmaceutical composition comprising a population of particles according to claim 24.



DEICOCO 1110 ---------

INLIEMNA LIUNAL SEAMUN REPURT

ir ional Application No PCT/GB 01/02922

A. CLASSI IPC 7	FICATION OF SUBJECT MATTER A61K9/16		
According to	o International Patent Classification (IPC) or to both national classification	n and IPC	
	SEARCHED		
Minimum do IPC 7	cumentation searched (classification system followed by classification $A61K$	symbols)	
Documental	ion searched other than minimum documentation to the extent that such	documents are included in the tields se	arched
	ata base consulted during the international search (name of data base at a, PAJ, EPO-Internal, CHEM ABS Data	and, where practical, search terms used)	
C. DOCUMI	ENTS CONSIDERED TO BE RELEVANT		
Calegory °	Citation of document, with indication, where appropriate, of the releva	int passages	Relevant to claim No.
Α	NL 7 501 406 A (PHARMACHIM) 10 August 1976 (1976-08-10) the whole document		1-25
A	WO 96 32095 A (ASTRA AKTIEBOLAGET) 17 October 1996 (1996-10-17) the whole document	1-25	
A,P	WO 00 38811 A (GLAXO) 6 July 2000 (2000-07-06) the whole document	1-25	
A,P	WO 00 44468 A (BRISTOL-MYERS SQUIBB) 3 August 2000 (2000-08-03) the whole document		1-25
			·
Funt	ner documents are listed in the continuation of box C.	Y Patent family members are listed i	n annex.
'A' docume consid 'E' earlier of filing of 'L' docume which citation 'O' docume other of the country of the cou	ent defining the general state of the art which is not ered to be of particular relevance document but published on or after the international ate ate int which may throw doubts on priority claim(s) or is cited to establish the publication date of another or or other special reason (as specified) ent referring to an oral disclosure, use, exhibition or neans ent published prior to the international filing date but	later document published after the inter or priority date and not in conflict with it cited to understand the principle or the invention document of particular relevance; the cleannot be considered novel or cannot involve an inventive step when the document of particular relevance; the cleannot be considered to involve an inventive an inventive an inventive and ocument is combined with one or moments, such combined with one or moments, such combination being obvious in the art.	he application but ory underlying the airmed invention be considered to surnent is taken alone airmed invention entive step when the re other such docusto a person skilled
		document member of the same patent f Date of mailing of the international sea	
	December 2001	14/12/2001	
Name and r	nailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2	Authorized officer	
European Patent Office, P.B. 5818 Patentiaan 2 NL - 2280 HV Rijswilk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl. Fax: (+31-70) 340-3016		Ventura Amat, A	

PCI	/GB	01/	02922
-----	-----	-----	-------

A	Publication date 10-08-1976 17-10-1996	NONE	Patent family member(s)	Publication date
		AU	694863 82	
. A	17-10-1996		694863 B2	
		AU CA CN EP JP NO NZ WO US ZA	5352496 A 2217062 A1 1186428 A 0820276 A1 11503448 T 974557 A 305515 A 9632095 A1 6221398 B1 9602596 A	30-07-1998 30-10-1996 17-10-1996 01-07-1998 28-01-1998 26-03-1999 02-10-1997 29-03-1999 17-10-1996 24-04-2001 14-10-1996
Α	.06-07-2000	AU BR EP WO NO	1877100 A 9916587 A 1144065 A1 0038811 A1 20013039 A	31-07-2000 25-09-2001 17-10-2001 06-07-2000 22-08-2001
A	03-08-2000	AU BR WO US	2513300 A 0007396 A 0044468 A1 6302958 B1	18-08-2000 30-10-2001 03-08-2000 16-10-2001
			A 06-07-2000 AU BR EP WO NO A 03-08-2000 AU BR WO	WO 9632095 A1 US 6221398 B1 ZA 9602596 A A 06-07-2000 AU 1877100 A BR 9916587 A EP 1144065 A1 WO 0038811 A1 NO 20013039 A A 03-08-2000 AU 2513300 A BR 0007396 A WO 0044468 A1

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

□ BLACK BORDERS
IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
FADED TEXT OR DRAWING
☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
☐ SKEWED/SLANTED IMAGES
☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
☐ GRAY SCALE DOCUMENTS
LINES OR MARKS ON ORIGINAL DOCUMENT
☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
Потивр.

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.